

60th Annual Scientific Session & Expo

E6

JACC April 5, 2011
Volume 57, Issue 14

CARDIAC ARRHYTHMIAS

REDUCTION IN LEFT ATRIAL VOLUMES WITH CARDIAC RESYNCHRONIZATION THERAPY AND THE RISK OF ATRIAL TACHYARRHYTHMIAS IN MADIT-CRT

ACC Oral Contributions

Ernest N. Morial Convention Center, Room 238

Monday, April 04, 2011, 8:00 a.m.-8:15 a.m.

Session Title: Atrial Tachyarrhythmias

Abstract Category: 26. Clinical Electrophysiology—Supraventricular Arrhythmias

Presentation Number: 906-3

Authors: *Andrew Brenyo, Alon Barsheshet, Arthur J. Moss, MD, Wojciech Zareba, Paul J. Wang, David T. Huang, Scott D. Solomon, Mark Link, Ilan Goldenberg, University of Rochester, Rochester, NY, Brigham and Women's Hospital - Harvard School of Public Health, Boston, MA*

Background: We hypothesized that reductions in left atrial volume (LAV) with a cardiac resynchronization therapy-defibrillator (CRT-D) will translate into subsequent reduction in the risk of atrial tachyarrhythmias.

Methods: Reductions of LAV at 1-year following CRT-D implantation (pre-specified as high- $\geq 20\%$ or low- $< 20\%$ LAV response) were related to the risk of subsequent atrial tachyarrhythmias (comprising atrial fibrillation, atrial flutter, or atrial tachycardia) among 1373 patients enrolled in MADIT-CRT.

Results: Mean LAV reductions among CRT-D and ICD-only patients were $28\% \pm 12\%$ vs. $10\% \pm 7\%$, respectively ($p < 0.001$). The cumulative probability of atrial tachyarrhythmias 3-years after assessment of echocardiographic response was lowest among those who showed a high LAV-response to CRT-D ($n=581$), and significantly higher among both low LAV-responders to CRT-D ($n=170$) and ICD-only patients ($n=622$; Figure). Congruently, multivariate analysis showed that high LAV-responders experienced a significant 57% ($p=0.029$) reduction in the risk of atrial tachyarrhythmias as compared with ICD-only patients, whereas the risk for the development of atrial tachyarrhythmias was not significantly different between low LAV-responders and ICD-only patients ($HR=1.05$; $p=0.89$).

Conclusions: In patients receiving CRT-D, favorable reverse atrial remodeling is associated with a significant reduction in the risk for the development of subsequent atrial tachyarrhythmias.

